Name:	Date:
Period:	Mr. Valentino

Unit 8 Test Topics

- Pythagorean Theorem
 - o Classifying triangles using Pythagorean theorem
- Trigonometry
 - o Sine, Cosine, Tangent
 - o Angle of Elevation and Depression
 - Law of Sines
 - o Law of Cosines
 - Cofunctions
- Special Right Triangles

Unit 9 Test Topics

- Properties of Parallelograms
- Properties of Rectangles
- Properties of Rhombuses
- Properties of Squares
- Properties of Trapezoids
 - o Midsegment, altitude problems
- Proofs with Quadrilaterals

Unit 10 Test Topics


- Slope
- Midpoint
- Distance
- Quadrilateral Proofs
- Triangle Proofs

Attached you will find 55 practice problems. This is not all-inclusive! You must study your notes, homework's and past tests!

- 1. In $\triangle ABC$, m $\angle A = 33$, a = 12, and b = 15. What is $m \angle B$ to the *nearest degree*?
 - 1) 41
 - 2) 43
 - 3) 44
 - 4) 48

- 2. In $\triangle ABC$, the complement of $\angle B$ is $\angle A$. Which statement is always true?
 - 1) $\tan \angle A = \tan \angle B$
 - 2) $\sin \angle A = \sin \angle B$
 - 3) $\cos \angle A = \tan \angle B$
 - sin ∠A = cos ∠B
- 3. In $\triangle FGH$, f = 6, g = 9, and $m \angle H = 57$. Which statement can be used to determine the numerical value of h?
 - 1) $h^2 = 6^2 + 9^2 2(9)(h)\cos 57^\circ$
- 3) $6^2 = 9^2 + h^2 2(9)(h)\cos 57^\circ$
- 2) $h^2 = 6^2 + 9^2 2(6)(9)\cos 57^\circ$
- 4) $9^2 = 6^2 + h^2 2(6)(h)\cos 57^\circ$
- 4. If $\sin 6A = \cos 9A$, then $m \angle A$ is equal to
 - 1) 6
 - 2) 36
 - 3) 45

Solve for x to the nearest degree.



- [A] 22
- [B] 68
- [C] 24
- [D] 66
- 6. ΔΧΥΖ is a right triangle with a right angle at Y. Which of the following is true?


[A]
$$\cos Z = \frac{XY}{XZ}$$
 [B] $\sin X = \frac{YZ}{XZ}$ [C] $\sin X = \frac{XY}{XZ}$ [D] $\tan X = \frac{XY}{ZY}$ [E] $\sin Z = \frac{YZ}{XZ}$

[B]
$$\sin X = \frac{YZ}{XZ}$$

[C]
$$\sin X = \frac{XY}{XZ}$$

[D]
$$\tan X = \frac{XY}{ZY}$$


- A large totem pole near Kalama, Washington, is 115 ft tall. On a particular day at noon it casts a 231 ft shadow. What is the sun's angle of elevation at that time?
- [A] 29.9° [B] 63.5° [C] 60.1°
- [D] 26.5°
- Use the Pythagorean theorem to solve for x.



- [A] √514
- [B] $\sqrt{208}$
- [C] √17
- [D] √64


- 9. In $\triangle ABC$, $\angle A$ is a right angle and $m \angle B = 45$. If AB = 17 feet, find AC.
 - [A] 14.722 ft
- [B] 29.445 ft
- [C] 17 ft
- [D] 24.042 ft

- 11. A 12-foot ladder is leaning against a building. The bottom of the ladder is 4 feet from the building. How far up the building is the top of the ladder? Round your answer to the nearest tenth.
- 13. A slide 3.5 m long makes an angle of 28° with the ground. How high is the top of the slide above the ground?
 - [A] 1.86 m
- [B] 1.64 m
- [C] 3.09 m
- [D] 1.71 m
- 15. A man 6 ft tall walks 75 ft from the base of a tree. He uses a protractor to measure the angle from his eye to the top of the tree. He finds it to be about 25°. Find the height of the tree to the nearest foot.
- 17. Solve for x to the nearest degree.



- [A] 58
- [B] 32
- [C] 62
- [D] 28

10. What is x to the nearest hundredth?




- [A] x = 11.44
- [B] x = 13.64
- [C] x = 17.61
- [D] x = 32.34
- 12. Which set of side lengths cannot form a right triangle?
 - [A] 12 mm, 16 mm, 20 mm
 - [B] 24 mm, 32 mm, 40 mm
 - [C] 6 mm, 8 mm, 10 mm
 - [D] 13 mm, 16 mm, 20 mm
- 14. A lookout spots a fire from a 36 meter tower. The angle of depression from the tower to the fire is 22 degrees. To the nearest meter, how far is the fire from the base of the tower?
- 16. Liola drives 12 km up a hill that is at a grade of 14°. What horizontal distance, to the nearest tenth of kilometer, has she covered?
 - [A] 13.7 km
- [B] 2.9 km
- [C] 3 km
- [D] 11.6 km
- 18. In $\triangle FUN$, f = 4, $m \angle F = 26$, and $m \angle N = 67$. Find the value of n to the *nearest integer*.

19. In $\triangle ABC$, a = 6, b = 12, and $m \angle C = 60$. What is the length of side c to the nearest integer?

- 1) 5
- 2) 10


- 3) 11
- 4) 20

20. Given the right triangle below, what is the length of the hypotenuse? Round your answer to the nearest tenth.




- [A] 181.0 cm
- [B] 13.5 cm
- [C] 4.4 cm
- [D] 19.0 cm

21. The diagram below shows the plans for a cell phone tower. A guy wire attached to the top of the tower makes an angle of 65 degrees with the ground. From a point on the ground 100 feet from the end of the guy wire, the angle of elevation to the top of the tower is 32 degrees. Find the height of the tower, to the nearest foot.




Find the measures of the numbered angles in the parallelogram.



- [A] $m \angle 1 = 28$; $m \angle 2 = 30$; $m \angle 3 = 122$
- [B] $m \angle 1 = 30$; $m \angle 2 = 28$; $m \angle 3 = 122$
- [C] $m \angle 1 = 15$; $m \angle 2 = 61$; $m \angle 3 = 150$
- [D] $m \angle 1 = 30$; $m \angle 2 = 15$; $m \angle 3 = 150$
- 25. The measures of the angles of a quadrilateral are x+15, 2x, x-45, and 2x-60. What type(s) of quadrilateral could this be?


 I. parallelogram II. rectangle III. trapezoid
 - [A] III only [B] I and II [C] II only
 - [D] I only [E] I and III
- 26. The cross section of an attic is in the shape of an isosceles trapezoid, as shown in the accompanying figure. If the height of the attic is 9 feet, <u>BC</u> = 12 feet, and <u>AD</u> = 28 feet, find the length of <u>AB</u> to the nearest foot.



23. In isosceles trapezoid JKLM, leg JK = 7x - 9, base KL = 5x + 3, and leg LM = 2x + 2. Find the value of x.


[A] 6 [B]
$$\frac{11}{5}$$
 [C] $-\frac{1}{3}$ [D] $-\frac{7}{5}$

24. Given ABCD is a rhombus, $m \angle ABC = 120$, and EB = 19. Find the length of \overline{AD} .



[A] 38 [B] 43 [C] 42 [D] 35

27. If ON = 8x - 2, LM = 7x + 7, NM = x - 4, and OL = 2y + 3, find the values of x and y for which LMNO must be a parallelogram.



[A]
$$x = 9$$
; $y = 1$ [B] $x = -\frac{1}{5}$; $y = -1$

[C]
$$x = 9$$
; $y = -1$ [D] $x = 5$; $y = -1$


- 28. A parallelogram must be a rectangle when its
 - diagonals are perpendicular
 - 2) diagonals are congruent
 - opposite sides are parallel
 - 4) opposite sides are congruent
- 30 A parallelogram is always a rectangle if
 - the diagonals are congruent
 - 2) the diagonals bisect each other
 - 3) the diagonals intersect at right angles
 - 4) the opposite angles are congruent
- 32. Which reason could be used to prove that a parallelogram is a rhombus?
 - Diagonals are congruent.
 - Opposite sides are parallel.
 - Diagonals are perpendicular.
 - Opposite angles are congruent.
- In quadrilateral ABCD, each diagonal bisects opposite angles. If m∠DAB = 70, then ABCD must be a
 - 1) rectangle
 - trapezoid
 - 3) rhombus
 - square
- 36. Which quadrilateral has diagonals that are always perpendicular bisectors of each other?
 - square
 - rectangle
 - trapezoid
 - parallelogram

- 29. Which quadrilateral has diagonals that always bisect its angles and also bisect each other?
 - rhombus
 - 2) rectangle
 - 3) parallelogram
 - isosceles trapezoid
- 31.

The diagonals of a quadrilateral are congruent but do not bisect each other. This quadrilateral is

- 1) an isosceles trapezoid
- a parallelogram
- a rectangle
- 4) a rhombus
- 33. Which quadrilateral does *not* always have congruent diagonals?
 - isosceles trapezoid
 - 2) rectangle
 - rhombus
 - square
- 35. In a certain quadrilateral, two opposite sides are parallel, and the other two opposite sides are not congruent. This quadrilateral could be a
 - rhombus
 - parallelogram
 - square
 - 4) trapezoid
- 37. Which statement is *false*?
 - 1) All parallelograms are quadrilaterals.
 - All rectangles are parallelograms.
 - All squares are rhombuses.
 - All rectangles are squares.


- Find the measure, to the nearest tenth, of the diagonal of a rectangle with dimensions 18 cm by 9 cm.
 - [A] 5.2 cm
- [B] 20.1 cm
- [C] 15.6 cm
- [D] 19 cm
- 40. Trapezoid TRAP, with median \overline{MQ} , is shown in the diagram below. Solve algebraically for x and y.



- In quadrilateral MNOP, ∠M ≅ ∠N.
 Quadrilateral MNOP could be a
 I. trapezoid. II. rhombus. III.
 parallelogram.
 - [A] I, II, or III
- [B] II or III

- [C] I or II
- [D] I only
- [E] III only

39. In rhombus *ABCD*, with diagonals \overline{AC} and \overline{DB} , AD = 10.




If the length of diagonal \overline{AC} is 12, what is the length of \overline{DB} ?

- 1) 8 2) 16 3) $\sqrt{44}$ 4) $\sqrt{136}$
- 41. <u>Isosceles trapezoid</u> ABCD has diagonals \overline{AC} and \overline{BD} . If AC = 5x + 13 and BD = 11x 5, what is the value of x?
 - 1) 28
 - 2) $10\frac{3}{4}$
 - 3) 3
 - 4) $\frac{1}{2}$
- 43. Select the geometric figure that possesses all of the following characteristics:
 - (1) quadrilateral
 - (2) diagonals equal
 - (3) opposite sides are parallel
 - [A] trapezoid
- [B] parallelogram
- [C] rhombus
- [D] rectangle


44. Given: AE ≅ EC, ED ≅ EB

 $AB \perp BC$, $AB \cong BC$

Prove: ABCD is a square




45. Given: *ABCD* is a rhombus. Prove: \triangle *BCA* \cong \triangle *DAC*



46. Given: $\triangle MJK \cong \triangle KLM$

Prove: MJKL is a parallelogram.



- 47. Which equation represents the perpendicular bisector of AB whose endpoints are A(8,2) and B(0,6)?
 - 1) y = 2x 4
 - 2) $y = -\frac{1}{2}x + 2$
 - 3) $y = -\frac{1}{2}x + 6$
 - 4) y = 2x 12

- 49. Line ℓ passes through the point (5,3) and is parallel to line k whose equation is 5x + y = 6. An equation of line ℓ is
 - 1) $y = \frac{1}{5}x + 2$
 - 2) y = -5x + 28
 - 3) $y = \frac{1}{5}x 2$
 - 4) y = -5x 28

- 51. The coordinates of A are (-9,2) and the coordinates of G are (3,14). What are the coordinates of the midpoint of AG?
 - 1) (-3,8)
 - 2) (-6,6)
 - 3) (-6, 16)
 - 4) (-21,-10)


- 48. What is an equation of the line that passes through the point (-2,5) and is perpendicular to the line whose equation is $y = \frac{1}{2}x + 5$?
 - 1) y = 2x + 1
 - 2) y = -2x + 1
 - 3) y = 2x + 9
 - 4) y = -2x 9

- 50. M is the midpoint of AB. If the coordinates of A are (-1,5) and the coordinates of M are (3,3), what are the coordinates of B?
 - 1) (1,4)
 - 2) (2,8)
 - 3) (7,1)
 - 4) (-5,7)

- 52. A line segment has endpoints (4,7) and (1,11). What is the length of the segment?
 - 1) 5
 - 2) 7
 - 3) 16
 - 4) 25

53. Given: Quadrilateral ABCD with A(-5, 0), B(1, -4), C(5, 2), D(-1, 6).

Prove: ABCD is a rectangle.



54. Quadrilateral *KATE* has vertices K(1,5), A(4,7), T(7,3), and E(1,-1).

a Prove that KATE is a trapezoid. [The use of the grid is optional.]

b Prove that KATE is not an isosceles trapezoid.

55. The coordinates of quadrilateral ABCD are A(-1,-5), B(8,2), C(11,13), and D(2,6). Using coordinate geometry, prove that quadrilateral ABCD is a rhombus. [The use of the grid is optional.]

